

## Dietary organic acids as non-antibiotic additives for aquaculture (in the Philippines)

Christian LÜCKSTÄDT

ADDCON Nordic AS, 3908 Porsgrunn, Norway  
christian.lueckstaedt@addcon.net

Growing awareness from consumers and producers of aquaculture species has resulted in a demand for responsible and sustainable aquaculture. The regulatory authorities in most exporting countries now focus on the misuse of antibiotic growth promoters (AGP) in aquaculture, while public attention has shifted towards sustainable production methods. As a result, alternatives to AGP have had to be found. Several feed additives, including acidifiers consisting of organic acids and their salts may be promising alternatives to the use of antibiotics in aquaculture feeds.

In recent trials in Southeast Asia, the inclusion of different organic acid salts in fish diets was tested for instance in tilapia, *Oreochromis niloticus*. One trial with tilapia fingerlings, weighing approximately 7 g, tested 5 different feeding treatments with 3 replicates each. The acidifier treatments (0.5%, 1.0% and 1.5%), containing Ca-formate, Ca-propionate, Ca-lactate, Ca-phosphate and citric acid, were tested against a negative control and a positive control containing an AGP (0.5% oxytetracycline). Fish were fed to satiation two times a day on a pelletized basal diet containing 31% crude protein for a period of 63 days. Although no statistically significant differences were observed, fish fed the 1.5% acidified diet had a numerically increased body weight gain (33.56 g) by more than 11% and 2% compared to the negative (30.18 g) and positive controls (32.98 g) respectively, while the 0.5% and 1.0% acidified diets had slightly poorer growth compared to the negative control. Tilapia which received the 1.5% acidifier dosage had furthermore an FCR improved by 3.5% compared to the negative control and by 1.4% compared to the AGP-treated fish.

Another experiment with male hybrid tilapia included 4 feed-treatment groups (negative control and 3 acidifier groups, containing 0.2%, 0.3% and 0.5% potassium diformate (Aquaform<sup>®</sup>), respectively). Fish were fed 6 times a day over an 85-day trial period. Beginning on day 10, all fish were orally challenged with *Vibrio anguillarum* ( $10^5$  CFU/d) once a day for 20 days. Significant improvements ( $P < 0.05$ ) could be monitored in all treated groups. The 0.2% application of potassium diformate in tilapia led to a significantly increased feed intake (8.6%), weight gain (18.6%) and a significantly improved feed conversion ratio (8.2%). Furthermore, mortality rates due to the *Vibrio* infection were significantly reduced.

With the results shown above (presented at World Aquaculture Society Meetings), it may be stated that the use of organic acid salts in tilapia aquaculture can improve the grow-out period in terms of performance and sustainability under Southeast Asian conditions.